

Resolving Domination Numbers of Family of Tree Graph

Regita Triani Mardiya

Departement of Mathematics Education
 University of Jember
 Jember, Indonesia
 diyaregita.99@gmail.com

Arika Indah Kristiana

Departement of Mathematics Education
 University of Jember
 Jember, Indonesia
 arika.fkip@unej.ac.id

Dafik

Departement of Mathematics Education
 University of Jember
 Jember, Indonesia
 d.dafik@unej.ac.id

Abstract: All graph in this paper are members of family of graph tree. Let G is a connected graph, for an ordered set $W = \{w_1, w_2, \dots, w_k\}$ of vertices and a vertex which is not element of W , then W is dominating set of graph G when the vertices that are not listed at W are vertices which are adjacent with W . The minimum cardinality of dominating set of graph G is called dominating numbers denoted $\gamma(G)$. If W and a vertex on graph G are connected each other, the metric representation of v which is element of W is the k -vector $r(v|W) = (d(v, w_1), d(v, w_2), \dots, d(v, w_k))$, where $d(x, y)$ represents distance between x and y . Then, W is resolving dominating set of graph G if the distance of all vertices is different respect to W . The minimum cardinality of resolving dominating set is called resolving domination numbers denoted $\gamma_r(G)$. In this paper we found the exact values of resolving dominating for firecracker graph, caterpillar graph and banana tree graph.

Keywords: Resolving Numbers, Domination Numbers, Resolving Domination Numbers, Family of Tree Graph.

1. INTRODUCTION (Heading 1)

Let $G(V, E)$ be a connected graph, then the resolving dominating set is a set of vertices on graph G that are members of dominating sets and resolving sets. This concept firstly was introduced by Bringham, et. al [1]. It combines two different concepts, which are concept of dominance and resolver. Concept of resolver is developed from the basic of minimum metrik concept which was studied by Sater in 1975 [2]. W is an ordered set of element of vertices on graph G where $W = \{w_1, w_2, \dots, w_k\}$, then W is dominating set of graph G when the vertices are not listed at W are vertices which is adjacent with W . The minimum cardinality of dominating set of graph G is called dominating numbers $\gamma(G)$. If W and a vertex on graph G are connected each other, the metric representation of v which is element of W is the k -vector $r(v|W) = (d(v, w_1), d(v, w_2), \dots, d(v, w_k))$, where $d(x, y)$ represents distance between x and y . Then, W is resolving dominating set of graph G if the distance of all vertices is different respect to W . The minimum cardinality

of dominating resolving set is called resolving domination numbers denoted $\gamma_r(G)$ [1]. The studies of domination resolving numbers, [3], [4], [5].

For illustration of placement of vertices which are element of domination resolving numbers is provided in Figure 1.




Figure 1: Vertices of resolving domination numbers of path, $\gamma_r(P_5) = 2$

Bringham, et. al [1] have found propotion of this topic, Propotion. For every graph G ,

$$\max\{\gamma(G), \dim(G)\} \leq \gamma_r(G) \leq \gamma(G) + \dim(G)$$

2. RESULT

Theorem 2.1. Resolving domination numbers of firecracker graph Fr_m^n for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) = nm$.

Proof. To prove that the resolving domination numbers of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) = nm$, it needs to be proven using the lower bound : $\gamma_r(Fr_m^n) \geq nm$ and the upper bound : $\gamma_r(Fr_m^n) \leq nm$

First, we prove the lower bound of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) \geq nm$. Assume that $\gamma_r(Fr_m^n) < nm$, we take $\gamma_r(Fr_m^n) = nm - 1$. Then we make possible placement of vertices of set W .

Possibility 1.

$$\begin{aligned} \text{Let } W = & \{y_i; 1 \leq i \leq n\} \cup \{z_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} \\ & - \{z_j^i; 1 \leq i \leq n, j = \{r\}, 1 \leq r \leq m\} \\ & - \{z_j^i; i = \{q\}, 1 \leq i \leq n, j = \{s\}, 1 \leq s \leq m\} \end{aligned}$$

From the construction above, we know that all vertices are dominated by vertices of element W , but there are two vertices that has the same representation of W . It shows contradiction. So we got lower bound of resolving domination numbers of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) \geq nm$.

Possibility 2.

$$\begin{aligned} \text{Let } W = & \{y_i; 1 \leq i \leq n\} - \{y_i; i = \{r\}, 1 \leq r \leq n\} \cup \\ & \{z_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} - \{z_j^i; 1 \leq i \leq n, \\ & j = \{s\}, 1 \leq s \leq m\} \end{aligned}$$

From the construction above, we know that each vertex has different representation of W , but there are two vertices that are not dominated by W . It shows contradiction. So we got lower bound of resolving domination numbers of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) \geq nm$.

Futhermore, we prove the upper bound of resolving dominating numbers of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) \leq nm$. Let

$W = \{y_i; 1 \leq i \leq n\} \cup \{z_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} - \{z_j^i; 1 \leq i \leq n, j = \{r\}, 1 \leq r \leq m\}$, so we got representation of vertices :

$$\begin{aligned} r(z_j^i | W) = & (2i - k, \underbrace{2i - k + 1, \dots, 2i - k + 1}_{m-1}, \underbrace{1, 2, \dots, 2}_{m-1}, \\ & \underbrace{h - i + 3, h - i + 4, \dots, h - i + 4}_{m-1}) \\ & \text{for : } 1 \leq i \leq n, 1 \leq k \leq i - 1, i + 1 \leq h \leq n \end{aligned}$$

$$\begin{aligned} r(x_i | W) = & (i - k + 1, \underbrace{i - k + 2, \dots, i - k + 2}_{m-1}, \underbrace{1, 2, \dots, 2}_{m-1}, \\ & \underbrace{h - i + 1, h - i + 2, \dots, h - i + 2}_{m-1}) \\ & \text{for : } 1 \leq i \leq n, 1 \leq k \leq i - 1, i + 1 \leq h \leq n \end{aligned}$$

From the construction above, we know that each vertex has different representation of W and dominated. So we got upper bound of resolving domination numbers of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) \leq nm$.

So, from those upper bound and lower bound we can conclude that the resolving domination numbers of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) = nm$.

Theorema 2.2. Resolving domination numbers of caterpillar graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) = nm$, it needs to be proven using the lower bound : $\gamma_r(Ct_m^n) \geq nm$ and the upper bound: $\gamma_r(Ct_m^n) \leq nm$.

First, we prove the lower bound of aterpillar graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) \geq nm$. Assume that $\gamma_r(Ct_m^n) < nm$, we take $\gamma_r(Ct_m^n) = nm - 1$. Then we make possible placement of vertices of set W .

Possibility 1.

$$\begin{aligned} \text{Let } W = & \{x_i; 1 \leq i \leq n\} \cup \{y_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} \\ & - \{y_j^i; 1 \leq i \leq m, j = \{r\}, 1 \leq r \leq m\} \end{aligned}$$

$$- \{y_j^i; i = \{q\}, 1 \leq q \leq m, j = \{s\}, 1 \leq s \leq m\}$$

From the construction above, we know that all vertices are dominated by vertices of element W , but there are two vertices that has the same representation of W . It shows contradiction. So we got lower bound of resolving domination numbers of caterpillar graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) \geq nm$.

Possibility 2.

$$\begin{aligned} \text{Let } W = & \{x_i; 1 \leq i \leq n\} - \{y_i; i = \{r\}, 1 \leq r \leq n\} \cup \\ & \{y_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} - \{y_j^i; 1 \leq i \leq n, \\ & j = \{s\}, 1 \leq s \leq m\} \end{aligned}$$

From the construction above, we know that each vertex has different representation of W , but there are two vertices that are not dominated by W . It shows contradiction. So we got lower bound of resolving domination numbers of caterpillar graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) \geq nm$.

Futhermore, we prove the upper bound of resolving dominating numbers of caterpillar graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) \leq nm$. Let $W = \{x_i; 1 \leq i \leq n\} \cup \{y_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} - \{y_j^i; 1 \leq i \leq n, j = \{r\}, 1 \leq r \leq m\}$, so we got representation of vertices :

$$\begin{aligned} r(y_j^i | W) = & \underbrace{(i - k + 1, i - k + 2, \dots, i - k + 2)}_{i-1}, \underbrace{1, 2, \dots, 2}_{m-1}, \\ & \underbrace{h + 1, h + 2, \dots, h + 2}_{m-1} \\ & \underbrace{}_{n-i} \\ \text{for } & 1 \leq i \leq n, 1 \leq k \leq i - 1, 1 \leq h \leq n - i \end{aligned}$$

From the construction above, we know that each vertex has different representation of W and dominated. So we got upper bound of resolving domination numbers of caterpillar graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) \leq nm$.

So, from those upper bound and lower bound we can conclude that the resolving domination numbers of firecracker graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) = nm$.

Theorema 2.3. Resolving domination numbers of banana tree graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) = nm$, it needs to be proven using the lower bound : $\gamma_r(B_m^n) \geq nm$ and the upper bound: $\gamma_r(B_m^n) \leq nm$.

First, we prove the lower bound of aterpillar graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) \geq nm$. Assume that $\gamma_r(B_m^n) < nm$, we take $\gamma_r(B_m^n) = nm - 1$. Then we make possible placement of vertices of set W .

Possibility 1.

$$\begin{aligned} \text{Let } W = & \{y_i; 1 \leq i \leq n\} \cup \{z_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} \\ & - \{z_j^i; 1 \leq i \leq n, j = \{r\}, 1 \leq r \leq m\} \\ & - \{z_j^i; i = q, 1 \leq q \leq n, j = \{s\}, 1 \leq s \leq m\} \end{aligned}$$

From the construction above, we know that all vertices are dominated by vertices of element W , but there are two vertices that has the same representation of W . It shows contradiction. So we got lower bound of resolving domination numbers of banana tree graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) \geq nm$.

Possibility 2.

$$\begin{aligned} \text{Let } W = & \{x_i; 1 \leq i \leq n\} - \{y_i; i = \{r\}, 1 \leq r \leq n\} \cup \\ & \{y_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} - \{y_j^i; 1 \leq i \leq n, \\ & j = \{s\}; 1 \leq s \leq m\} \end{aligned}$$

From the construction above, we know that each vertex has different representation of W , but there are two vertices that are not dominated by W . It shows contradiction. So we got lower bound of resolving domination numbers of caterpillar graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) \geq nm$.

Futhermore, we prove the upper bound of resolving dominating numbers of firecracker graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) \leq nm$. Let

$$\begin{aligned} W = & \{y_i; 1 \leq i \leq n\} \cup \{z_j^i; 1 \leq i \leq n, 1 \leq j \leq m\} \\ & - \{z_j^i; 1 \leq i \leq n, j = \{r\}, 1 \leq r \leq m\}, \text{ so we got } \\ & \text{representation of vertices :} \end{aligned}$$

$$r(z_j^i | W) = (\underbrace{3, \dots, 3}_{i-1}, 1, \underbrace{3, \dots, 3}_{n-1}, \underbrace{4, \dots, 4}_{(i-1)(m-1)}, \underbrace{2, \dots, 2}_{m-1}, \underbrace{4, \dots, 4}_{(n-i)(m-1)})$$

for : $1 \leq i \leq n$

$$r(x_i | W) = (\underbrace{1, \dots, 1}_n, \underbrace{2, \dots, 2}_{n(m-1)})$$

From the construction above, we know that each vertex has different representation of W and dominated. So we got upper bound of resolving domination numbers of banana tree graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) \leq nm$.

So, from those upper bound and lower bound we can conclude that the resolving domination numbers of banana tree graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) = nm$.

3. CONCLUSION

In this paper we have been studied about resolving domination numbers of family of tree graph. We have been concluded the exact value of firecracker graph Fr_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(Fr_m^n) = nm$, the resolving domination numbers of firecracker graph Ct_m^n , for $n \geq 1$ and $m \geq 2$ is $\gamma_r(Ct_m^n) = nm$, and the resolving domination numbers of banana tree graph B_m^n , for $n \geq 2$ and $m \geq 1$ is $\gamma_r(B_m^n) = nm$.

4. ACKNOWLEDGMENT (HEADING 5)

We gratefully acknowledge the support from CGANT University of Jember Indonesia of year 2019.

5. REFERENCES

- [1] Bringham, R.C., Gary C., Ronald D., Dutton, and Ping Z. (2003). Resolving Domination in Graphs, Journal Mathematica Bohemica, vol. 128, no. 1, pp. 25-36.
- [2] Slater, P.J. (1975). Maximin Facility Location, Journal of Research of the National Bureau of Standart - B. Mathematical Science, vol. 79B, no. 3-4.
- [3] Slater, P.J. (1987). Domintion and Location in Acyclic Graphs, Netwoks, vol. 17, no. 55-64.
- [4] Hernando, C., M. Mora, I.M. Pelayo. (2018). Resolving Dominationg Partitions in Graphs.
- [5] Stephen, S. Bharati R., Cyriac G., and Albert W. (2015). Resolving-Power Dominating Sets. Applied Mathematics and Computation. Vol. 256, no. 778-785.