

Efficacy of Insecticides against Infestation of Brinjal Fruit Borer, *Leucinodes orbonalis* (Pyralidae: Lepidoptera) in District Awaran, Balochistan

¹Ameer Uddin, ²Muhammad Yousuf, ³Waqar Ali Lasi, ⁴Muhammad Naeem, ⁵Abdul Ghafoor, ⁶Dr. Rehan Fareed, ⁷Mitha Khan, ⁸Siraj Ahmed, ⁹Allah Bakhsh, ¹⁰Khalil Ahmed, ¹¹Zia Ul Haq

¹ Acting Entomologist/Deputy Director/Horticulturist, Agriculture Research, District Awaran, 01900, Balochistan, Pakistan

Main Author Gmail address: uddinameer302@gmail.com

² Acting Horticulturist, Directorate of Agriculture Research, Panjgoor, 93000, Pakistan

³ Director, Directorate of Agriculture Research Wayaro Farm District Lasbela at Uthal, 90150, Pakistan

⁴ Director, Directorate of Agriculture Research (Dates) at Turbat, 92600, Pakistan

⁵ Acting Deputy Director (Statistics), Directorate of Agriculture Research Wayaro Farm District Lasbela at Uthal, 90150, Pakistan

⁶ Soil Chemist/Deputy Director, Directorate of Agriculture Research Wayaro Farm District Lasbela at Uthal, 90150, Pakistan

⁷ Acting Entomologist/Deputy Director, Directorate General of Agriculture Research Institute (ARI), Sariab, Quetta, 87550, Pakistan

⁸ Acting Soil Chemist, Directorate of Agriculture Research (Dates) at Turbat, 92600, Pakistan

⁹ Agriculture Officer, Agriculture Extension Department, District Awaran, 01900, Balochistan, Pakistan

¹⁰ Acting Plant Pathologist/Deputy Director, Directorate of Agriculture Research (Medicinal Plant) Kalat, 88300, Pakistan

¹¹ Acting Plant Pathologist, Directorate of Agriculture Research Fodder, ARI, Sariab, Quetta, 87550, Pakistan

Abstract: The experiment research trial to study the efficacy of different insecticides against Brinjal fruit borer was studied in the Agriculture Research District Awaran, Balochistan in the year 2022 under the supervision of Agriculture Research Awaran. The insecticides tested were Match, Proclaim, Tracer, Talstar, Steward and Larvin were compared with a control. Results showed that Proclaim insecticide was most effective against brinjal fruit borer and resulted in lower infestation (40.1%) followed by Larvin (40.43%), whereas Steward offered moderate control (41.31%) of brinjal fruit borers. The study recommended the use of Proclaim insecticide is most effective control of brinjal fruit borer in district Awaran and field condition.

Keywords: Brinjal fruit borer, Efficacy, Insecticides, Agriculture Research, District Awaran, Balochistan.

Introduction:

Brinjal fruit borer, *Leucinodes orbonalis* Guenee (Pyralidae: Lepidoptera) is found throughout the tropics in Asia and Africa and is a minor pest in America. It is monophagous and very important pest of brinjal, however other plants belonging to family Solanaceae are reported to be hosts of this pest. They include tomato (*Lycopersicon esculentum*), potato (*Solanum tuberosum*) and turkey berry (*S. torvum*). It is an internal borer which damages the tender shoots and fruits. Attack of this pest causes considerable damage to brinjal crop each year, affecting the quality and yield of the crop. The Larvae of this pest cause 12-16% damage to shoots and 20-60% damage to fruits. The pest is very active during rainy and summer season and often causes more than 90% damage in Bangladesh and up to 95% in India. It is also reported that the infestation of fruit borer causes reduction in Vitamin C content to an extent of 68 % in the infested fruits. Soon after hatching from eggs,

young caterpillars search for and bore into tender shoots near growing points into flower buds or into the fruits. Caterpillars prefer fruits over other plant parts. Larvae go through at least five instars and there are reports of the existence of six larval instars. Larval period lasts from 12 to 15 days in the summer and up to 22 days in winter. Climatic conditions are important in the life cycle of the borer. As temperature increases and humidity decreases fecundity increases and duration of life cycle decreases. Within one hour after hatching, the larvae bore into the nearest tender shoot, flower or fruit, they plug the entrance hole with excreta. In young plants, caterpillar bores midrib of large leaves. As a result, the affected leaves may drop off. Larvae feeding inside shoots result in wilting of young shoots. Presence of wilted shoots in a brinjal field is a symptom of damage by this pest. The damaged shoots ultimately wither and drop off. This reduces plant growth, which in turn, reduces fruit number and size. New shoots can arise but this delays

crop maturity and the newly formed shoots are also subjected to larval damage. Larval feeding in flowers is a relatively rare occurrence resulting in failure to form fruits from damaged flowers. Larval feeding inside the fruit results in destruction of fruits tissue. The feeding tunnels are often clogged with frass. This makes even slightly damaged fruit unfit for marketing. The yield loss varies from season to season and from location to location. Damage to fruits

Materials and Methods:

Research Area: The present study was conducted at Agriculture Research District Awaran under the supervision of Agriculture Research Awaran in the year 2022 to study the efficacy of six insecticides against *L. orbonalis* under field conditions. Brinjal was sown under field conditions in which spacing was provided by keeping plant to plant distance of 20 cm and row to row distance of 40 cm. Six insecticides that is Match, Proclaim, Tracer, Talstar, Steward and Larvin were tested along with a control. The field

Statistical Analysis:

The values of mean percent damage were then statistically analyzed by analysis of variance. Least significance differences (LSD) was determined at the probability level of $P < 0.05$ to decide the significance of individual treatment effect.

Results:

The results are summarized in the illustrates to study the efficacy of six insecticides against *L. orbonalis* under field conditions. The data collected on damaged fruits before spraying indicated no significant difference among the percent fruit damage in all the treatments (Table 1). Data recorded before the spray application showed a mean percent damage of 69.53, 57.3, 51.63, 57.93, 51.63, 56.53 and 52.16% on different treatments. Control treatment showed a percent fruit damage of 59.73%. Data recorded after 3rd days of application of insecticides revealed lowest percentage of fruit damage in Proclaim which reduced the percent damage from 57.3% to 39.1% followed by Talstar 43.3%, Steward 44.43%, Tracer 46.33%, Larvin 46.40% and Match 48.66% (Table 1). The percent fruit damage in treated plots of Tracer, Larvin, Steward, Talstar were not significantly different from each other but significantly different from Proclaim and control. During 2022 in district Awaran, the minimum pest infestation was recorded in Steward treated plots followed by Proclaim 3rd days after application of insecticides. The maximum pest infestation was recorded in control plot (Table 2).

particularly in autumn, is very severe and the whole crop can be destroyed. Based on the mentioned facts the present study was initiated to check the efficacy of different novel insecticides for controlling this devastating insect pest. The objective of this experiment research trial was to determine the to study the efficacy of different insecticides against Brinjal fruit borer was studied in the Agriculture Research District Awaran, Balochistan and Pakistan.

experiment was laid out in randomized complete block design (RCBD) and each treatment was replicated three times. Spraying was done by using air compressing knapsack sprayer. The observations on count of damage fruits were recorded on three randomly selected plants per treatment. First count was done one day before insecticide application and post treatment counts were made after 3, 6, 9, 12 and 15 days. In order to assess the percent fruit damage, infested fruits on three randomly selected plants were counted as against total number of available fruits on the observed plants.

Data on percent fruit damage after 6th days of application indicated no significant difference among all the treatments. The percent fruit damage in Proclaim was on par with Talstar, Steward and Larvin, while they were significantly different from the control. Match, Tracer and control treatments were not significantly different from each other. Lowest percentage of damage fruits was found in Proclaim i.e. 43.66% and Talstar 43.10% followed by Steward i.e. 43.2%, Larvin 44.03%, Tracer 45.56%. The highest percentage of damaged fruits was found in plots treated with Match i.e. 49.36% (Table 1). There was significant difference in insecticide efficacy six days after application of insecticides in the year, 2022 in the district Awaran. The minimum pest infestation was recorded in Steward treated plots (Table 2). The percent fruit infestation after 6th day of application was not significantly different from each other in treated but significantly different from the control. The highest percent fruit infestation was recorded in plots treated with Tracer i.e. 44.46% followed by Match and Talstar (42.40 and 41.16%), whereas the lowest (40.20%) damage was record in Proclaim followed by Larvin and Steward i.e., 40.60 and 40.56% (Table 1). Steward was the most effective insecticide showing the minimum pest infestation six days after spray in 2022 in district Awaran. The data recorded after 9th day of spraying revealed lowest percent fruit infestation in Proclaim i.e. 26.93% which was followed by Match, Larvin, Steward and Talstar i.e. 32.66, 30.90, 36.6 and 39.60%. The highest (43.16%)

damage was recorded in Tracer. The percent fruit infestation in Proclaim treated plots was significantly different from Talstar, Tracer, Steward and control treatments, while it was on par with Match and Larvin. There was no significant difference in percent fruit damage in Tracer, Talstar and Steward treated plots (Table 1). The 9th days post treatment data showed that Steward was most effective insecticide in second spray compared with other insecticides (Table 2). After 12th day of application table revealed lowest percent fruit infestation of 29.23% was recorded in Proclaim followed by Larvin, Match, Steward and Tracer i.e. 29.43, 32.20, and 43.03%, respectively. The highest (43.03%) fruit infestation was recorded in Talstar. No significant difference was recorded in percent fruit damage among Proclaim, Match, Larvin and Steward treatments, while they were significantly different

from Tracer, Talstar and control. Percent damage in Tracer treated plots was significantly different from rest of the treatments as well as from the control after 12th day of insecticide application (Table 1). Tracer showed maximum residual toxicity and minimum pest infestation 15th days post treatment compared with other insecticides (Table 2). Overall mean of the data indicated a significant difference among the treatments and control. The lowest (40.1%) fruit infestation was recorded in Proclaim. The highest percent fruit infestation of 58.15% was recorded in control. The result showed that Proclaim was found to be the most effective in suppressing the fruit infestation by brinjal fruit borer, while Steward offered moderate control of brinjal fruit borer. The overall mean of second spray showed that Steward was the most effective insecticide followed by Match and Proclaim in reducing pest infestation.

Table 1: Percent infestation of brinjal fruit borer after application of insecticides during 2022 under field conditions in district Awaran

TREATMENTS	BEFORE SPRAY	TIME INTERVALS					MEANS
		3 rd Day	6 th Day	9 th Day	12 th Day	15 th Day	
MATCH	69.53	48.66b	49.36a b	42.40b	32.66de	29.23c	45.31b
PROCLAIM	57.30	39.10d	43.06b	40.20b	26.93c	32.10c	40.1c
TRACER	57.93	43.33bc	45.56ab	44.46b	43.16b	43.03b	46.75b
TALSTAR	56.53	43.40c	43.10b	41.16b	39.60bc	56.03a	46.64b
STEWARD	52.16	44.43c	43.20b	40.60b	36.60de	32.13c	41.31c
LARVIN	51.63	46.40bc	44.03b	40.56b	30.90de	29.03c	40.43c
CONTROL	59.73	58.7a	53.33a	60.90a	54.93a	61.30a	58.15a
LSD 5%		3.06	7.95	8.01	6.53	9.17	1.76

Means within a column followed by different letters are significantly different from each other at 5% level of significance (LSD Test)

Table 2: Percent infestation of brinjal fruit borer after application of insecticides during 2022 under field conditions in district Awaran

TREATMENTS	BEFORE SPRAY	TIME INTERVALS					MEANS
		3 rd Day	6 th Day	9 rd Day	12 th Day	15 th Day	
MATCH	60.45	50.81ab	49.54bc	45.53 _b	46.56 _b c	38.77bc	46.24bc
PROCLAIM	50.30	42.09ab	45.53bc	56.59 _a b	39.44 _b c	51.93b	47.11bc
TRACER	53.93	52.17ab	67.37abc	48.54 _a b	50.81b	35.89c	50.95bc
TALSTAR	50.53	53.47ab	79.38a	49.54 _a b	42.07 _b c	38.29bc	52.55b
STEWARD	45.16	31.27b	42.09c	42.09b	33.91c	50.95bc	40.06c
LARVIN	49.63	51.65ab	60.03abc	50.81 _a b	43.89 _b c	39.42bc	49.16bc
CONTROL	69.53	60.01a	69.07ab	65.03a	73.98a	67.82a	67.18a
LSD 5%		24.95	25.36	22.02	16.72	15.71	12.07

Means within a column followed by different letters are significantly different from each other at 5% level of significance (LSD Test)

Discussion:

The present observations on the effectiveness of Proclaim are in conformity with those of Anil and Sharma (2010). These results also coincide with Kalawate and Dethé (2012) who stated that Proclaim and Match are efficient in reducing the population and the subsequent damage caused by brinjal fruit borer. Insecticides are currently the main method of control for *Leucinodes orbonalis*. Contact insecticides are the most commonly used and show varying degrees of an efficacy against the pest. Deltamethrin and endosulfan were the most effective insecticides used in South Asia. In field experiments, triazophos and methomyl were applied when more than 20 % of brinjal fruits were infested. The highest fruit yields were obtained with triazophos. Babu *et al.*, (2002) found that the most commonly used insecticides in the field were carbaryl, followed by endosulfan, carbofuran and cypermethrin in India. A combination of cypermethrin/deltamethrin and triazophos/endosulfan sometimes combines with cartap hydrochloride and diflubenzuron gave higher yields than non-treated plots,

cypermethrin/deltamethrin mixtures were the most effective. Studies made by Kalawate and Dethé (2012) on the bio efficacy of Match and Proclaim in comparison to Larvin and self formulated neem extract for two cropping seasons against brinjal fruit and shoot borer indicated that Match was found to be the most effective against *Leucinodes orbonalis* although the corresponding yield recorded in Larvin, Control was higher but it was not significantly different than that noticed in Match and Proclaim. Anil and Sharma (2010) observed the bio efficacy of insecticides against brinjal shoot and fruit borer. The result showed that in terms of shoot infestation, Proclaim (0.002%), endosulfan (0.005%), novaluron (0.01%) and lambda-cyhalothrin (0.004%) were found superior. The total number of drooping shoots was minimum in Proclaim followed by endosulfan and novaluron as compared to Match, deltamethrin and *Bacillus thuringiensis*, in terms of reduction in fruit infestation, Proclaim (0.002%) was highly effective followed by endosulfan (0.05%) agrospray oil T (0.2%) and Match (0.002%). However, cost benefit ratio was highest in agrospray oil T (0.2%)

followed by lambda-cyhalothrin (0.004%) endosulfan (0.05%) and deltamethrin (0.0028%). Rosaiah (2001) sprayed the plant products against brinjal shoot and fruit borer, neemazol at 0.5 per cent was significantly superior in reducing shoot and fruit damage and contributed to maximum yield. The next best treatments were neemazol 0.25 percent, NSKE (neem seed kernel and jatropha were poor in their effectiveness against brinjal shoot and fruit borer. Patel *et al.*, (2001) found that one spray of monocrotophos 0.04 percent at 9 weeks after transplanting followed by three sprays of spark (triazophos + deltamethrin) 0.036 per cent at 11, 13 and 15 weeks after transplanting effectively managed the brinjal fruit borer and gave significantly higher fruit yield (238.8 q/ha) percent on fruits, respectively.

References:

1. Alam MZ. Insect pest of vegetables and their control in Bangladesh. Agril. Inf. Serv. Dacca. Bangladesh, 1970, 132.
2. Maurel AM, Noriel LM, Esguerra NM. Life history and behavior of eggplant fruit borer. *Annal. Trop. Res* 1982; 4(3):178.
3. Ali MI, Rahman MS. Field evaluation if wilt disease and shoot and fruit borer attack of different cultivators of brinjal. *Bangladesh J Agric Sci* 1980; 7(2):193-194.
4. Kalloo. Solanaceous crops. In *Vegetable Breeding Raton, Florida* 1988; 2:520-570.
5. Naresh JS, Malik VS, Balan JS, Khokhar KS. A New record of trathala sp, a larval endoparasite attacking brinjal fruit borer, *Leucinodes orbonalis* Guenee. *Bull. Entomol. New Delhi* 1986; 27(1):74.
6. Hemi MA. Effect of borer attach on the vitamin 'C' content of brinjal. *Pakistan J. Health* 1955; 4:223-224.
7. Atwal AS. *Agricultural pests of Indian and Southern Asia*. New Delhi: Kalyani publishers, 1976, 529.
8. Kumar P, Johnsen S, Life cycle studies on fruit and shoot borer (*Leucinodes orbonalis*) and natural enemies of insect-pests of eggplant (*Solanum melongena*). *Journal of Applied Biology* 2000; 10(2):178-184.
9. Bhutani DK, Jotwani MR. *Insects of vegetables in New Delhi: Periodical expert book Agency*, 1984, 356.
10. Anil, Sharma PC. Bioefficacy of insecticides against *Leucinodes orbonalis* on brinjal. *J. Environmental biology* 2010; 4(2):31-32.
11. Kalawate A, Dethe MD. Bioefficacy study of biorational insecticides on brinjal. *J. Biopesticide* 2012; 5(1):75-80.
12. Thanki KV, Patel JR. Efficacy and economics of certain insecticides used for the control of pest complex on brinjal. *Gujrat Agric U Res J* 1991; 16(2):47-51.
13. Radhika S, Reddy KD, Subbarathnam GV. Management of brinjal shoot and fruit borer *Leucinodes orbonalis* (Guenee) with insecticides. *J. Research Angra* 1997; 25(3):10-14.
14. Babu BS, Varaprasad KS, Anitha KP, Rao SK. Chakarbarti, Chandurkar PS. Brinjal shoot and fruit borer (*Leucinodes orbonalis*) status in coastal districts of Orissa. *Resources management in plants protection during twenty first century, Hyderabad, India* 2002; 2:121-123.
15. Kumar BA, Rao BN, Sriramulu M. Studies on the efficacy of certain insecticides and their mixtures against shoot and fruit borer, *Leucinodes orbonalis* on brinjal in India. *J. Plt. Prot* 2000; 28(1):25-28.
16. Babu BS, Varaprasad KS, Anitha KP, Rao SK. Chakarbarti, Chandurkar PS. Brinjal shoot and fruit borer (*Leucinodes orbonalis*) status in coastal

Conclusions:

The present experiment research trial finding results demonstrates and showed that Proclaim insecticide was most effective in suppressing the pest population as lowest mean percent fruit infestation was recorded after its application i.e. 40.1% followed by Larvin in which percent fruit infestation was recorded 40.43%. The Steward and Match gave moderate control of brinjal fruit borer in which the mean percent fruit infestation was 41.31% and 45.31%. It is suggested that Proclaim insecticide should be used for controlling this pest in district Awaran. However, in case of unavailability of this insecticide Larvin may be applied.

- districts of Orissa. Resources management in plants protection during twenty first century, Hyderabad, India 2002; 2:121-123.
17. Kumar BA, Rao BN, Sriramulu M. Studies on the efficacy of certain insecticides and their mixtures against shoot and fruit borer, *Leucinodes orbonalis* on brinjal in India. J. Plt. Prot 2000; 28(1):25-28.
18. Biradar AP, Taggelli RG, Patil DR, Balikai RA. Evaluation of cypermethrin 3 EC + quinalphos 20 EC against brinjal shoot and fruit borer. Karnataka J. Agric. Sci 2001; 142(2):369-372.
- Patel JJ, Patel BH, Maghodia AB, Bhalala MK. Integration Tolerant Variety and insecticides application for the management of jassids and fruit borer infesting brinjal. J. Pestology 2001; 25:56.