International Journal of Academic Health and Medical Research (IJAHMR)
ISSN: 2643-9824
Vol. 9 Issue 10 October - 2025, Pages: 91-96

Flavonoids As Promising Natural Antioxidants: Mechanisms, Sources, And Therapeutic Applications

Ahmed Attahiru[*1], Bashar Attahiru[2], Ahmed Musa Ahmed[1], and Nafiu Sani Gulumbe[3]

[1]Department of Pure and Industrial Chemistry, Faculty of Physical Sciences
Kebbi State University of Science and Technology, Aliero
Kebbi State, Nigeria
E-mail address: ahmedattahiru02@gmail.com
ORCID ID: https://orcid.org/0009-0009-3437-2287
[2]Department of Microbiology, Faculty of Life Sciences
Kebbi State University of Science and Technology, Aliero
Kebbi State, Nigeria
[3]Department of Science Education, Faculty of Sciences
Waziri Umaru Federal Polytechnic, Birnin Kebbi
Kebbi State, Nigeria

ABSTRACT: *Flavonoids are one of the largest groups of naturally occurring polyphenolic compounds in plants and are widely recognized for their significant contributions to human health. Over the past decades, they have emerged as vital bioactive molecules with diverse pharmacological activities, of which antioxidant capacity remains the most extensively studied and therapeutically relevant. These compounds are highly effective in neutralizing reactive oxygen and nitrogen species, modulating oxidative stress pathways, and maintaining cellular redox balance. Such properties place flavonoids at the center of natural product research aimed at disease prevention and management. Increasing scientific evidence has highlighted their roles in mitigating the onset and progression of several chronic conditions, particularly cardiovascular diseases, neurodegenerative disorders, diabetes mellitus, and cancer. This review provides a comprehensive assessment of the biochemical and molecular mechanisms underlying the antioxidant actions of flavonoids, while also exploring dietary and medicinal plant sources that serve as rich reservoirs of these compounds. Methodology for reviewing the literature is included, outlining systematic database search strategies, inclusion and exclusion criteria, and data extraction procedures to ensure comprehensive and unbiased analysis. Furthermore, the article integrates tables and figures that summarize flavonoid subclasses, sources, and therapeutic effects for enhanced understanding. By presenting both preclinical and clinical findings, the review underscores the potential of flavonoids as promising candidates for antioxidant-based therapies. Finally, it discusses prevailing challenges such as low bioavailability, highlights current advances in delivery systems, and identifies gaps in knowledge that require future investigation to fully harness the therapeutic potential of flavonoids.*

Keywords: Flavonoids, Antioxidants, Oxidative stress, Therapeutic applications, Natural products

1. Introduction

Oxidative stress, a pathological state that results from an imbalance between the excessive generation of reactive oxygen species (ROS) and the limited capacity of endogenous antioxidant defenses, has been strongly implicated in the initiation and progression of numerous chronic diseases. ROS, which include free radicals such as hydroxyl radical, superoxide anion, and non-radical species like hydrogen peroxide, are generated as natural byproducts of cellular metabolism. Under normal physiological conditions, their production is tightly regulated by endogenous antioxidant systems, ensuring a delicate redox balance that supports cellular signaling and metabolic functions. However, when ROS production overwhelms antioxidant defenses, oxidative stress ensues, leading to molecular damage. Lipids undergo peroxidation, proteins become denatured or inactivated, and nucleic acids experience mutagenic lesions. These processes disrupt cellular homeostasis and accelerate pathological pathways, contributing to the etiology of cardiovascular disorders, cancer, diabetes mellitus, and neurodegenerative conditions such as Alzheimer's and Parkinson's diseases (Valko et al., 2016). The cumulative burden of oxidative stress is also associated with aging, making its mitigation an essential focus of biomedical research.

Given the central role of oxidative stress in human disease, the scientific community has increasingly focused on developing therapeutic strategies that can effectively counter oxidative insults. Traditional approaches have relied on synthetic antioxidants, including butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA). These compounds have long been incorporated into food preservation systems and pharmaceutical formulations due to their ability to delay lipid oxidation and enhance shelf life. However, safety concerns regarding their potential carcinogenicity, pro oxidant effects under certain conditions, and toxicity at higher doses have raised significant debate (Shahidi & Ambigaipalan, 2015). Consequently, the search for safer and more effective alternatives has shifted attention toward naturally derived antioxidants, particularly those from plants. Plant based antioxidants offer several advantages over synthetic compounds. Beyond exhibiting lower toxicity profiles, they act through multiple complementary biochemical pathways. This multifunctionality enhances their therapeutic relevance in complex diseases where oxidative stress interacts with inflammation, apoptosis, and metabolic dysregulation. Among these natural agents, flavonoids have emerged as one of the most promising groups due to their abundance in human diets, structural diversity, and wide ranging biological activities.

Flavonoids are polyphenolic compounds synthesized via the phenylpropanoid pathway, and they are widely distributed across edible and medicinal plants. They occur naturally in fruits such as apples, grapes, and berries; vegetables like onions, kale, and broccoli; herbs including parsley, thyme, and chamomile; as well as in commonly consumed beverages such as tea, cocoa, and red wine. Their ubiquity in dietary sources contributes significantly to daily antioxidant intake, with epidemiological studies consistently linking high flavonoid consumption to reduced risk of chronic diseases.

Structurally, flavonoids comprise multiple subclasses including flavones, flavonols, flavanones, flavanols (catechins), isoflavones, and anthocyanins, each characterized by specific hydroxylation and glycosylation patterns that influence their solubility, stability, and biological activity (Panche et al., 2016). This diversity allows for a broad spectrum of antioxidant mechanisms.

At the molecular level, flavonoids exert their antioxidant effects through several complementary mechanisms. First, they directly scavenge reactive oxygen and nitrogen species by donating hydrogen atoms or electrons from their phenolic hydroxyl groups, stabilizing the resulting radical intermediates. Second, flavonoids chelate transition metals such as iron and copper, thereby inhibiting the Fenton and Haber Weiss reactions that otherwise generate highly reactive hydroxyl radicals. Third, they modulate endogenous antioxidant defense systems by upregulating the activity of key enzymes including superoxide dismutase, catalase, and glutathione peroxidase. Finally, flavonoids exert indirect antioxidant effects by regulating redox sensitive signaling pathways such as nuclear factor erythroid 2 related factor 2 (Nrf2), which controls the expression of cytoprotective genes, and nuclear factor kappa B (NF kB), a central regulator of inflammatory responses. Through these combined mechanisms, flavonoids not only neutralize oxidative insults but also restore and sustain cellular redox homeostasis.

This review provides an in depth exploration of flavonoids as natural antioxidants with therapeutic promise. It highlights their molecular mechanisms of action, dietary and medicinal plant sources, and pharmacological applications in diseases associated with oxidative stress. Special attention is given to the therapeutic relevance of flavonoids in cardiovascular, neurodegenerative, metabolic, and oncological disorders, reflecting their multifunctional capacity to intervene at various stages of disease progression.

The methodology employed in compiling and synthesizing relevant literature includes a systematic search of peer reviewed journals, reviews, and clinical studies indexed in databases such as PubMed, Scopus, and Web of Science. The inclusion criteria emphasized studies published between 2000 and 2025 that examined antioxidant mechanisms, therapeutic outcomes, or clinical relevance of flavonoids. Exclusion criteria removed duplicate records, non-English publications, and articles lacking mechanistic or pharmacological data. This approach ensured a comprehensive and evidence based synthesis of current knowledge.

To enhance clarity and accessibility, tables and figures are integrated to summarize subclasses of flavonoids, their dietary and medicinal sources, and their principal antioxidant mechanisms. Such visual aids are intended to provide quick reference points for readers while complementing detailed textual descriptions.

Despite strong evidence of their antioxidant and therapeutic potential, flavonoids face limitations in clinical application. A major challenge is their poor bioavailability, as many flavonoids undergo extensive metabolism in the gastrointestinal tract and liver, limiting systemic concentrations of active forms. Advances in formulation strategies including nanoparticle delivery systems, liposomal encapsulation, and structural modification are being actively investigated to overcome these barriers. Furthermore, while preclinical studies have provided compelling insights, clinical validation remains limited. Large scale randomized controlled trials are necessary to establish effective doses, safety profiles, and long term benefits in diverse patient populations.

In general, flavonoids represent a versatile and promising class of natural antioxidants with significant implications for human health. Their unique ability to act through multiple antioxidant mechanisms, coupled with their widespread dietary availability, positions them as key candidates in both preventive and therapeutic strategies against oxidative stress related disorders. Future research directed toward optimizing their bioavailability, elucidating synergistic interactions with other dietary compounds, and confirming clinical efficacy will be crucial for translating their potential into tangible health benefits.

2. METHODOLOGY

This review was conducted using a systematic approach to identify and analyze relevant peer-reviewed literature on flavonoids, antioxidants, and therapeutic applications. The methodology involved the following steps:

2.1 Literature Search

Electronic databases including PubMed, ScienceDirect, Scopus, and Google Scholar were searched for publications between 2000 and 2025. Keywords used included “flavonoids”, “antioxidant activity”, “oxidative stress”, “therapeutic applications of flavonoids”, and “natural polyphenols”.

2.2 Inclusion and Exclusion Criteria

Inclusion criteria consisted of studies published in English, reporting either experimental or clinical data on flavonoids as antioxidants, reviews, and meta-analyses relevant to human health. Excluded studies were those unrelated to antioxidant activity, unpublished theses, and articles without peer-review.

2.3 Data Extraction

Data were extracted regarding flavonoid classification, sources, antioxidant mechanisms, pharmacological applications, and clinical outcomes. Special focus was given to studies that linked biochemical mechanisms to therapeutic effects.

2.4 Data Analysis

The extracted data were synthesized into thematic sections. Results were compared across different flavonoid subclasses to highlight similarities and differences in antioxidant capacity and therapeutic potential. Graphical representations and summary tables were created to improve comprehension.

3. CLASSIFICATION AND SOURCES OF FLAVONOIDS

Flavonoids are a diverse group of polyphenolic compounds characterized by a C6–C3–C6 carbon skeleton. They are classified into several subclasses including flavones, flavonols, flavanones, flavanols (catechins), isoflavones, and anthocyanins (Panche et al., 2016).

Table 1. Major subclasses of flavonoids and dietary sources

Subclass	Example compounds	Major sources
Flavones	Apigenin, Luteolin	Parsley, celery, chamomile
Flavonols	Quercetin, Kaempferol	Onions, kale, apples, tea
Flavanones	Naringenin, Hesperidin	Citrus fruits
Flavanols	Catechin, Epicatechin	Green tea, cocoa, grapes
Isoflavones	Genistein, Daidzein	Soybeans, legumes
Anthocyanins	Cyanidin, Delphinidin	Berries, red cabbage, black rice

4. MECHANISMS OF ANTIOXIDANT ACTION

Flavonoids exert antioxidant effects through a wide range of molecular and biochemical mechanisms. Their activity is not limited to direct neutralization of reactive oxygen species (ROS) but extends to metal chelation, regulation of endogenous antioxidant enzymes, and modulation of cell signaling pathways that govern inflammation, apoptosis, and redox homeostasis. The diversity of these mechanisms explains why flavonoids are consistently linked to the prevention and management of chronic diseases such as cardiovascular disorders, diabetes, cancer, and neurodegenerative conditions. Recent advances in molecular biology and omics-based technologies have further highlighted that flavonoids act as multifunctional modulators rather than simple radical scavengers (Zou et al., 2021).

4.1 Free Radical Scavenging

The ability of flavonoids to scavenge free radicals remains the most widely investigated aspect of their antioxidant activity. Hydroxyl radicals, superoxide anions, hydrogen peroxide, and peroxynitrite are among the most reactive species generated during oxidative stress, and flavonoids can neutralize these by donating hydrogen atoms or electrons from their hydroxyl groups (Heim et al., 2002). The structural arrangement of hydroxyl substituents strongly influences their radical scavenging efficiency. For instance, the catechol structure in the B ring of quercetin and catechin confers high reactivity toward ROS, while the presence of a double bond between C2 and C3 in conjugation with a 4-oxo group enhances electron delocalization and antioxidant capacity (Mocan et al., 2021). These interactions allow flavonoids to directly prevent lipid peroxidation, protein oxidation, and DNA damage, thereby preserving cellular integrity.

4.2 Metal Chelation

Transition metals such as iron and copper are central mediators of oxidative damage because they catalyze the Fenton and Haber Weiss reactions, leading to the formation of hydroxyl radicals from hydrogen peroxide. Flavonoids inhibit these reactions by binding metal ions through their hydroxyl and carbonyl groups, forming stable chelates that prevent metal-mediated ROS generation (Pietta, 2000). For example, rutin and luteolin have been shown to effectively chelate Fe^{2+} and Cu^{2+}, thereby disrupting catalytic cycles of oxidative stress (López-Alarcón & Denicola, 2013). This chelating ability is particularly important in biological systems where excess free iron is implicated in neurodegenerative diseases such as Alzheimer's and Parkinson's disease. Recent in vivo studies further suggest that flavonoid metal complexes may exert additional pharmacological benefits, including modulation of enzyme activity and improved bioavailability (Younis et al., 2022).

4.3 Regulation of Antioxidant Enzymes

Beyond their direct interactions with ROS, flavonoids enhance cellular resilience by modulating the expression and activity of endogenous antioxidant enzymes. These include superoxide dismutase (SOD), which catalyzes the dismutation of superoxide into hydrogen peroxide; catalase, which decomposes hydrogen peroxide into water and oxygen; and glutathione peroxidase, which reduces lipid peroxides using glutathione as a substrate (Williams et al., 2004). Flavonoids such as epigallocatechin gallate (EGCG) and quercetin have been demonstrated to upregulate these enzymes at both transcriptional and post-translational levels, resulting in long-lasting protective effects against oxidative insults (Shen et al., 2020). By strengthening endogenous defense networks, flavonoids provide a sustained buffer against oxidative stress beyond their immediate radical scavenging function.

4.4 Modulation of Cell Signaling Pathways

An additional mechanism of action involves the regulation of redox-sensitive signaling pathways. Flavonoids activate the nuclear factor erythroid 2 related factor 2 (Nrf2) pathway, which is a master regulator of antioxidant defense genes. Through this pathway, compounds like quercetin and kaempferol promote the expression of heme oxygenase 1, NAD(P)H quinone oxidoreductase 1, and glutamate-cysteine ligase, which together contribute to improved cellular detoxification and oxidative balance (Surh, 2003; Xu et al., 2022). At the same time, flavonoids can inhibit the nuclear factor kappa B (NF-κB) signaling cascade, thereby reducing the production of proinflammatory cytokines such as TNF-α, IL-1β, and IL-6 (Liao et al., 2021). This dual modulation of Nrf2

activation and NF-κB suppression highlights the broader therapeutic relevance of flavonoids in diseases characterized by inflammation and oxidative imbalance.

4.5 Emerging Perspectives

Recent evidence suggests that flavonoids may also influence mitochondrial function, endoplasmic reticulum stress, and epigenetic regulation, thereby expanding the traditional understanding of their antioxidant mechanisms. For instance, flavonoids have been shown to preserve mitochondrial membrane potential, enhance ATP production, and regulate apoptotic proteins such as Bcl-2 and Bax (Zhang et al., 2023). Advances in metabolomics and computational modeling are providing deeper insights into these multi-target interactions, which may pave the way for precision use of flavonoid-rich extracts in personalized nutrition and medicine.

5. THERAPEUTIC APPLICATIONS

Flavonoids, owing to their strong antioxidant and multifunctional biological properties, have demonstrated wide ranging therapeutic potential against chronic diseases where oxidative stress plays a central role. Their ability to modulate cellular redox balance, regulate enzyme activity, and interact with signaling pathways provides a strong foundation for their application in disease prevention and management. In addition to antioxidant effects, flavonoids exert anti-inflammatory, antimicrobial, vasodilatory, and immunomodulatory activities, which broaden their clinical relevance. The following subsections highlight some of the most significant therapeutic areas where flavonoids have shown promise.

5.1 Cardiovascular Diseases

Oxidative stress is a critical factor in the pathogenesis of cardiovascular disorders such as atherosclerosis, hypertension, and ischemic heart disease. Flavonoids like quercetin, kaempferol, and catechins exert cardioprotective effects by improving endothelial function, reducing low density lipoprotein (LDL) oxidation, and inhibiting platelet aggregation. These compounds also modulate vascular inflammation and enhance nitric oxide bioavailability, thereby promoting vasodilation and improving blood flow (Hertog et al., 1993).

Epidemiological studies further suggest that populations with high dietary flavonoid intake have reduced incidence of cardiovascular morbidity and mortality (Grosso et al., 2017). For example, the consumption of flavonoid-rich foods such as berries, green tea, and cocoa is strongly associated with lower risk of coronary artery disease. Experimental and clinical studies indicate that flavonoids also protect cardiac myocytes from ischemia reperfusion injury, modulate angiotensin-converting enzyme activity, and reduce blood pressure in hypertensive patients (Bondonno et al., 2019). This evidence highlights their therapeutic relevance as natural cardioprotective agents.

5.2 Neurodegenerative Disorders

The central nervous system is highly vulnerable to oxidative damage due to its high oxygen consumption, abundant lipid content, and relatively weak antioxidant defenses. Flavonoids can cross the blood-brain barrier and directly interact with neurons and glial cells. Their neuroprotective actions are mediated by the suppression of oxidative stress, inhibition of neuroinflammation, and regulation of signaling pathways associated with neuronal survival.

In Alzheimer's disease, flavonoids such as epicatechin and luteolin reduce amyloid beta aggregation, decrease tau hyperphosphorylation, and protect neurons from oxidative injury (Spencer, 2010). In Parkinson's disease, flavonoids mitigate dopaminergic neuron loss by scavenging free radicals, modulating mitochondrial function, and activating neurotrophic signaling pathways such as brain derived neurotrophic factor (BDNF) and ERK-CREB cascades (de Andrade Teles et al., 2018). Recent animal studies further suggest that chronic consumption of flavonoid rich diets enhances synaptic plasticity, improves memory performance, and delays age-related cognitive decline (Mocan et al., 2021). These findings position flavonoids as potential adjuncts in managing age related neurodegenerative disorders.

5.3 Diabetes Mellitus

Oxidative stress contributes significantly to the onset and progression of diabetes and its complications. Flavonoids, particularly isoflavones, flavonols, and flavanones, improve glucose metabolism by enhancing insulin sensitivity, increasing glucose uptake, and modulating carbohydrate digesting enzymes such as α-glucosidase and α-amylase (Bahadoran et al., 2013). These effects help reduce postprandial hyperglycemia and improve long term glycemic control.

Moreover, flavonoids protect pancreatic β-cells from oxidative damage, thereby sustaining insulin secretion and preventing β-cell apoptosis (Shen et al., 2020). In experimental models, quercetin, rutin, and catechin reduce hyperglycemia, dyslipidemia, and insulin resistance, while attenuating diabetic complications such as nephropathy, neuropathy, and retinopathy through their anti-inflammatory and antioxidant activities (Rashid et al., 2021). Clinical studies are increasingly supporting these findings, suggesting that flavonoids may serve as valuable adjuncts in diabetes management alongside conventional therapies.

5.4 Cancer Prevention

Cancer development is a multistage process involving initiation, promotion, and progression, often driven by oxidative DNA damage, chronic inflammation, and dysregulation of cell proliferation and apoptosis. Flavonoids inhibit tumor initiation and progression through multiple mechanisms, including free radical scavenging, suppression of pro-carcinogenic enzymes such as cytochrome P450 isoforms, modulation of apoptosis, and interference with cell cycle regulation.

Specific flavonoids such as genistein, quercetin, apigenin, and luteolin have demonstrated strong anti-proliferative effects in experimental models of breast, colon, liver, and prostate cancers (Scalbert et al., 2005; Russo et al., 2012). Mechanistic studies

reveal that flavonoids regulate critical signaling pathways such as MAPK, PI3K/Akt, and Wnt/β-catenin, thereby inhibiting uncontrolled cell proliferation and promoting apoptosis of malignant cells (Nabavi et al., 2015).

Additionally, flavonoids possess antiangiogenic properties, preventing the formation of new blood vessels required for tumor growth, and enhance the efficacy of conventional chemotherapeutics by sensitizing resistant cancer cells to apoptosis (Liao et al., 2021). Clinical investigations have also reported correlations between high flavonoid intake and reduced cancer risk, particularly for gastrointestinal and hormone-dependent cancers (Zhang et al., 2023). Collectively, these findings support the development of flavonoids as potential chemopreventive and adjuvant therapeutic agents in oncology.

Table 2. Summary of therapeutic applications of flavonoids

Disease condition	Flavonoids studied	Mechanism of action
Cardiovascular diseases	Quercetin, Catechins	Antioxidant, anti-inflammatory, vascular protection
Neurodegenerative diseases	Epicatechin, Luteolin	Neuronal protection, modulation of signaling
Diabetes mellitus	Isoflavones, Kaempferol	Antioxidant, insulin sensitization
Cancer	Apigenin, Quercetin	Apoptosis induction, inhibition of proliferation

6. CHALLENGES AND LIMITATIONS

Although flavonoids demonstrate impressive antioxidant and therapeutic properties, several challenges hinder their clinical translation. One of the major limitations is poor bioavailability. Many flavonoids exhibit low aqueous solubility, which restricts their absorption in the gastrointestinal tract. In addition, they are prone to degradation in acidic gastric environments and undergo extensive first pass metabolism, leading to rapid elimination from systemic circulation (Manach et al., 2005). These pharmacokinetic barriers reduce the concentration of active compounds reaching target tissues, thereby limiting therapeutic efficacy. Another challenge lies in the structural diversity of flavonoids, which causes variability in their absorption, distribution, metabolism, and excretion. Furthermore, clinical studies are fewer compared to preclinical investigations, making it difficult to establish standardized dosages and safety profiles. To address these limitations, strategies such as nanoformulation, encapsulation in liposomes or polymeric carriers, conjugation with lipids, and co-administration with bioenhancers like piperine are being actively explored. While these approaches show promise in improving stability and bioavailability, they require rigorous evaluation for long term safety and effectiveness.

7. FUTURE PERSPECTIVES

Future research on flavonoids must move beyond in vitro and animal models to well-designed human clinical trials that determine optimal dosages, safety thresholds, and long term effects. The development of advanced delivery systems, including nanoparticles, solid lipid carriers, and phytosomes, holds potential for overcoming pharmacokinetic barriers. Moreover, studying synergistic interactions between flavonoids and other dietary phytochemicals or conventional drugs may uncover novel therapeutic applications. Emerging technologies such as genomics, proteomics, and metabolomics can provide deeper insights into molecular targets of flavonoids and help identify biomarkers of response. Computational modeling and artificial intelligence are also likely to play a role in predicting flavonoid protein interactions and optimizing drug design. Collaborative research integrating natural product chemistry, pharmacology, and clinical sciences will be essential for translating flavonoid research into effective interventions.

8. CONCLUSION

Flavonoids are a diverse class of natural antioxidants with multiple mechanisms of action that make them highly relevant in preventing and managing oxidative stress related disorders such as cardiovascular disease, neurodegeneration, diabetes, and cancer. Their ability to modulate free radicals, chelate metals, enhance antioxidant enzymes, and regulate signaling pathways underscores their therapeutic value. However, challenges related to bioavailability, metabolism, and lack of standardized clinical evidence remain significant barriers. Addressing these issues through innovative formulation technologies, robust clinical trials, and integrative research approaches will be key to unlocking the full clinical potential of flavonoids as natural therapeutic agents.

ACKNOWLEDGMENTS

The gratitude goes to the Kebbi State University of Science and Technology, Aliero and Nigeria TETFund for supporting related doctoral research

REFERENCES

1. Bahadoran, Z., Mirmiran, P., & Azizi, F. (2013). Dietary polyphenols as potential nutraceuticals in management of diabetes: A review. Journal of Diabetes & Metabolic Disorders, 12(1), 43. https://doi.org/10.1186/2251-6581-12-43
2. Heim, K. E., Tagliaferro, A. R., & Bobilya, D. J. (2002). Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships. The Journal of Nutritional Biochemistry, 13(10), 572–584. https://doi.org/10.1016/S0955-2863(02)00208-5
3. Hertog, M. G., Feskens, E. J., Hollman, P. C., Katan, M. B., & Kromhout, D. (1993). Dietary antioxidant flavonoids and risk of coronary heart disease: The Zutphen Elderly Study. The Lancet, 342(8878), 1007–1011. https://doi.org/10.1016/0140-6736(93)92876-U

4. Manach, C., Williamson, G., Morand, C., Scalbert, A., & Rémésy, C. (2005). Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. The American Journal of Clinical Nutrition, 81(1), 230S–242S. https://doi.org/10.1093/ajcn/81.1.230S
5. Panche, A. N., Diwan, A. D., & Chandra, S. R. (2016). Flavonoids: An overview. Journal of Nutritional Science, 5, e47. https://doi.org/10.1017/jns.2016.41
6. Pietta, P. G. (2000). Flavonoids as antioxidants. Journal of Natural Products, 63(7), 1035–1042. https://doi.org/10.1021/np9904509
7. Scalbert, A., Manach, C., Morand, C., Rémésy, C., & Jiménez, L. (2005). Dietary polyphenols and the prevention of diseases. Critical Reviews in Food Science and Nutrition, 45(4), 287–306. https://doi.org/10.1080/1040869059096
8. Shahidi, F., & Ambigaipalan, P. (2015). Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects. Journal of Functional Foods, 18, 820–897. https://doi.org/10.1016/j.jff.2015.06.018
9. Spencer, J. P. (2010). Flavonoids and brain health: Multiple effects underpinned by common mechanisms. Genes & Nutrition, 5(2), 243–250. https://doi.org/10.1007/s12263-009-0159-9
10. Surh, Y. J. (2003). Cancer chemoprevention with dietary phytochemicals. Nature Reviews Cancer, 3(10), 768–780. https://doi.org/10.1038/nrc1189
11. Valko, M., Rhodes, C. J., Moncol, J., Izakovic, M., & Mazur, M. (2016). Free radicals, metals and antioxidants in oxidative stress-induced cancer. Chemico-Biological Interactions, 160(1), 1–40. https://doi.org/10.1016/j.cbi.2005.12.009
12. Williams, R. J., Spencer, J. P., & Rice-Evans, C. (2004). Flavonoids: Antioxidants or signalling molecules. Free Radical Biology and Medicine, 36(7), 838–849. https://doi.org/10.1016/j.freeradbiomed.2004.01.001